

BRIEF REPORT

The Influence of the Dissociative Subtype of Posttraumatic Stress Disorder on Treatment Efficacy in Female Veterans and Active Duty Service Members

Erika J. Wolf

National Center for PTSD at VA Boston Healthcare System,
Boston, Massachusetts, and Boston University School of
Medicine

Carole A. Lunney

National Center for PTSD, White River Junction, Vermont

Paula P. Schnurr

National Center for PTSD, White River Junction, Vermont, and Geisel School of Medicine at Dartmouth

Objective: A dissociative subtype of posttraumatic stress disorder (PTSD) was recently added to the *Diagnostic and Statistical Manual of Mental Disorders* (5th ed.; American Psychiatric Association, 2013) and is thought to be associated with poor PTSD treatment response. **Method:** We used latent growth curve modeling to examine data from a randomized controlled trial of prolonged exposure and present-centered therapy for PTSD in a sample of 284 female veterans and active duty service members with PTSD to test the association between the dissociative subtype and treatment response. **Results:** Individuals with the dissociative subtype (defined using latent profile analysis) had a flatter slope ($p = .008$) compared with those with high PTSD symptoms and no dissociation, such that the former group showed, on average, a 9.75 (95% confidence interval $[-16.94, -2.57]$) lesser decrease in PTSD severity scores on the Clinician Administered PTSD Scale (Blake et al., 1995) over the course of the trial. However, this effect was small in magnitude. Dissociative symptoms decreased markedly among those with the subtype, though neither treatment explicitly addressed such symptoms. There were no differences as a function of treatment type. **Conclusions:** Results raise doubt about the common clinical perception that exposure therapy is not effective or appropriate for individuals who have PTSD and dissociation, and provide empirical support for the use of exposure treatment for individuals with the dissociative subtype of PTSD.

What is the public health significance of this article?

This study found that female veterans and active duty service members with the dissociative subtype of posttraumatic stress disorder (PTSD) did not respond as well to PTSD treatment with prolonged exposure or present-centered therapy as did those without the subtype. However, both PTSD and dissociation symptoms did improve markedly in the dissociative group, suggesting that the dissociative subtype is not a contraindication for the use of empirically supported treatments for PTSD.

Keywords: dissociative subtype, PTSD, latent growth curve, treatment response

Supplemental materials: <http://dx.doi.org/10.1037/ccp0000036.supp>

This article was published Online First July 13, 2015.

Erika J. Wolf, National Center for PTSD at VA Boston Healthcare System, Boston, Massachusetts, and Department of Psychiatry, Boston University School of Medicine; Carole A. Lunney, National Center for PTSD, White River Junction, Vermont; Paula P. Schnurr, National Center for PTSD, White River Junction, Vermont, and Department of Psychiatry, Geisel School of Medicine at Dartmouth.

The data reported in this article were drawn from Grant CSP #494 from the VA Cooperative Studies Program with support from the Department of

Defense for CSP #494. This work was also supported by a Career Development Award to Erika J. Wolf from the United States (U.S.) Department of Veterans Affairs, Clinical Sciences Research and Development Program. The contents of this article do not represent the views of the U.S. Department of Veterans Affairs, the Department of Defense, or the United States Government.

Correspondence concerning this article should be addressed to Paula P. Schnurr, National Center for PTSD (116D), VA Medical Center, White River Junction, VT 05009. E-mail: paula.schnurr@dartmouth.edu

The dissociative subtype (DS) of posttraumatic stress disorder (PTSD) was newly added to the fifth edition of the *Diagnostic and Statistical Manual of Mental Disorders (DSM-5; American Psychiatric Association, 2013)*, and is defined as meeting full criteria for PTSD and showing comorbid “persistent or recurrent” symptoms of derealization (i.e., experiencing the world as unreal or dreamlike) and/or depersonalization (i.e., feeling as if one’s physical body is disconnected from the self). Psychometric research, primarily using latent profile analysis (LPA), has provided consistent evidence across populations that approximately 15% to 30% of those with PTSD manifest the DS (e.g., Armour, Elklit, Lauterbach, & Elhai, 2014; Stein et al., 2013; Steuwe, Lanius, & Frewen, 2012; Wolf, Lunney, et al., 2012; Wolf, Miller, et al., 2012).

The inclusion of the DS in the *DSM-5 (American Psychiatric Association, 2013)* may yield new insight into the pathophysiology and treatment of PTSD. Dissociation is thought to negatively influence the efficacy of exposure-based PTSD therapies (Lanius et al., 2010), such as prolonged exposure (PE), because it may prevent activation of the fear network (i.e., processing of trauma-related emotions; van Minnen, Harned, Zoellner, & Mills, 2012). Further, research suggests that clinicians consider dissociation a contraindication for exposure therapy for PTSD (Becker, Zayfert, & Anderson, 2004). Despite this, no studies have found that dissociation negatively influences response to exposure therapy (Hagenaars, van Minnen, & Hoogduin, 2010; Halvorsen, Stenmark, Neuner, & Nordahl, 2014; Price, Kearns, Houry, & Rothbaum, 2014).

Dissociation is also thought to impair the ability to generate, attend to, and retain restructured trauma-related cognitions (a key component of cognitive therapies for PTSD) and to limit access to trauma-related memories (necessary for narrative therapies). But, to date, the data support only subtle effects of dissociation on response to cognitive and narrative therapies. Specifically, Resick, Suvak, Johnides, Mitchell, and Iverson (2012) reported no overall effect of dissociation on PTSD symptom decline in response to cognitive processing therapy (CPT), however, individuals with high levels of dissociation undergoing the full course of CPT showed faster PTSD symptom improvement relative to those undergoing only cognitive therapy. Similarly, Cloitre, Petkova, Wang, and Lu Lassell (2012) reported no overall effect of baseline dissociation on change in PTSD in a randomized controlled trial of Skills Training in Affective and Interpersonal Regulation (STAIR) and Narrative Story Telling (NST). However, participants with high posttreatment dissociation who completed the combined STAIR/NST arm showed greater PTSD symptom decline at 3- and 6-month follow-ups than did individuals with high posttreatment dissociation in the other treatment conditions. In both Resick et al. and Cloitre et al., the differences that emerged in PTSD response were within the groups with relatively high levels of dissociation and not between those with high versus low levels of dissociation.

Aims and Hypotheses

The primary aim of this study was to examine whether the DS was associated with differential PTSD symptom response to treatment in a sample of female service members and veterans. A second aim was to evaluate how dissociative symptoms changed among those with the DS in response to PTSD treatment. We

evaluated these aims in a data set that we have previously employed to test the latent structure of dissociation and PTSD. Specifically, using data from a large, randomized clinical trial of PE versus present-centered therapy (PCT) for the treatment of PTSD among military women (Schnurr et al., 2007), we previously conducted LPAs of the 17 *DSM-IV (American Psychiatric Association, 1994)* PTSD symptoms and four items indexing derealization and depersonalization, as measured at the baseline assessment Wolf, Lunney, et al. (2012). Approximately 30% of the sample was assigned to the dissociative class. To extend this, we retained class assignment (the moderate-PTSD, high-PTSD, or high-PTSD and high-dissociation groups) as a predictor of PTSD symptoms and symptom change using latent growth curve modeling. We hypothesized that the DS would be associated with poorer response to treatment as defined by less PTSD symptom improvement relative to the moderate- and high-PTSD groups, and that dissociative symptoms would improve among those with the DS.

Method

Participants and Measures

Participants were 284 female veterans ($n = 277$) and active-duty service members ($n = 7$) with PTSD who participated in a multisite ($n = 12$) randomized clinical trial of PE for PTSD (Schnurr et al., 2007). Mean age of participants was 44.79 years ($SD = 9.44$, range = 22 to 78). Almost half (45.4%, $n = 129$) self-identified as a non-White minority. The majority of the sample had been exposed to sexual assault (93.0%). Data were available from 235 participants at immediate posttreatment, 232 participants at 3-month follow-up, and 229 participants at 6-month follow-up. Participants were not removed from the study if they missed an assessment. PTSD was assessed using the Clinician-Administered PTSD Scale (CAPS; Blake et al., 1995). Dissociation was measured using four self-report derealization and depersonalization items from the Dissociation scale of the Trauma Symptom Inventory (Briere, 1995).¹

Procedure

Institutional review boards approved the protocol. A master’s- or doctoral-level assessor, blinded to condition, performed assessments before and after treatment and at 3- and 6-month follow-ups. All sessions were videotaped and reviewed by supervisors who provided weekly phone supervision; group calls with master supervisors were employed to maintain treatment fidelity. Participants were randomly assigned to 10 weekly 90-min sessions of PE ($n = 141$) or PCT ($n = 143$). PE included psychoeducation, imaginal exposure, homework, and discussion of thoughts and feelings related to exposures. PCT focused on current life problems as manifestations of PTSD and on general daily difficulties, accomplishments made during therapy, and future plans. There were no significant trauma or demographic differences among the

¹ Additional participant descriptive characteristics, and details about the measures and procedures, including inclusion/exclusion criteria and response rates and attrition, are provided in Schnurr et al. (2007).

Table 1

Observed Mean CAPS Severity Scores (and SDs) by Group and Treatment Type at Each Assessment

Class	Baseline		Month 0		Month 3		Month 6	
	PCT	PE	PCT	PE	PCT	PE	PCT	PE
Moderate PTSD	64.75 (9.20)	64.44 (10.06)	51.25 (19.69)	36.72 (25.16)	47.59 (20.61)	37.00 (23.03)	45.63 (23.01)	36.91 (22.88)
High-PTSD	91.03 (12.00)	89.21 (9.00)	64.26 (25.27)	67.93 (27.26)	62.33 (27.38)	60.90 (27.70)	61.06 (27.53)	63.57 (28.11)
High-PTSD & dissociation	83.98 (15.29)	86.21 (17.65)	70.62 (23.53)	66.83 (27.72)	65.25 (25.62)	65.69 (26.47)	65.00 (30.13)	64.40 (29.49)

Note. CAPS = Clinician-Administered PTSD Scale; PCT = present-centered therapy; PE = prolonged exposure; PTSD = posttraumatic stress disorder. All values are *M* (*SD*).

treatment groups and no site or therapist treatment effects (Schnurr et al., 2007).

Data Analyses

We created two dummy-coded variables to reflect LPA class assignment with the dissociative class as the reference group (moderate-PTSD vs. the high-PTSD and dissociative class; high-PTSD vs. the high-PTSD and dissociative class) and specified them as predictors of the latent intercept and slope factors in a latent growth curve model. As the shape of change over time was not a focus of this research, and because we did not expect the shape of change to be equivalent across all time points, we fixed the first and last slope loadings (Baseline and Month 6) and freely estimated those for Months 0 and 3. To obtain estimated means at Months 0 and 3, we fixed the Baseline and Month 0 (and, separately, the baseline and Month 3) slope loadings in follow-up analyses. We also examined Treatment Type \times Class as a predictor of the slope and intercept factors. Analyses were conducted separately for the CAPS scores and the derealization/depersonalization item means. We examined whether the groups differed in terms of loss of PTSD diagnosis over time using logistic regression and in the amount of treatment received as a function of class (and Class \times Treatment Type) using chi square and ANOVAs. For all analyses, we compared the dissociative class with the moderate- and high-PTSD classes, though the primary comparison of interest was that between the dissociative and the high-PTSD classes because differences between these groups can be attributed to dissociation. In contrast, the dissociative class differed from the moderate-PTSD group on both dissociation and PTSD severity, making it impossible to determine the source of treatment trajectory differences. Latent variable analyses were conducted in Mplus 7.0 (Muthén & Muthén, 1998–2012) using maximum likelihood estimation. Missing data were modeled directly (maximum amount of missingness was 20.1%).

Results

Descriptive statistics for each class at each time point as a function of treatment type are listed in Table 1. An initial latent growth curve model that included Treatment Type \times Class interaction terms as predictors of the intercept and slope factors fit the data well, $\chi^2(13, N = 284) = 10.33, p = .67$, root mean square error of approximation (RMSEA) $< .001$, 90% confidence interval (CI) for RMSEA [$< .001, .05$], comparative fit index (CFI) = 1.0, Tucker-Lewis Index (TLI) = 1.01, standardized root mean square residual (SRMR) = .02, but revealed no significant interactions

between Treatment Type and Class (smallest $p = .29$). Therefore, growth models collapsed across treatment type.

The second model, which included the effects of group membership on the slope and intercept factors, yielded a good fitting model, $\chi^2(7, N = 284) = 3.78, p = .80$, RMSEA [$< .001, .05$] 90% CI for RMSEA $< .001 = .05$, CFI = 1.0, TLI = 1.01, SRMR = .01, with the moderate- and high-PTSD groups, relative to the dissociative group, predicting the latent intercept. The high-PTSD group's baseline CAPS score was, on average, 5.05 points greater than that of the dissociative class, 95% CI [1.29, 8.81], $p = .009$, and the moderate-PTSD group's baseline CAPS score was, on average, 20.49 points below that of the dissociative class, 95% CI [$-23.89, -17.10$], $p < .001$. There was also an effect of the high-PTSD class (vs. the dissociative class) on the slope factor: On average, CAPS scores decreased 9.75 points more from the first to the last assessments in the high-PTSD class relative to the dissociative class, 95% CI [$-16.94, -2.57$], $p = .008$ (see Figure 1).² There was no slope effect comparing the moderate-PTSD group with the dissociative one (unstandardized [unst] $\gamma = -3.69, p = .26$), nor was the baseline CAPS score associated with the rate of change (unst $\gamma = 24.74, p = .76$). The mean baseline CAPS score for the high-PTSD and dissociative class was estimated as 85.09, 95% CI [82.49, 87.69], $p < .001$, and the mean rate of change for this group from the first to last assessment was -18.71 , 95% CI [$-23.75, -13.68$], $p < .001$. The residual variance for the slope was significant ($p < .001$), suggesting variables other than class assignment contributed to individual differences in rate of change; residual variance in the intercept was not significant ($p = .08$).

Logistic regressions revealed an omnibus significant group difference in the percentage of women who no longer met criteria for PTSD at any posttreatment assessment, $\chi^2(2, N = 235) = 18.19, p = .0001$ (see Table 2). However, follow-up pairwise testing showed that this was a function of differences between the moderate-versus high-PTSD group, $\chi^2(1, N = 235) = 10.34, p = .0013$, and versus the dissociative group, $\chi^2(1, n = 235) = 12.20, p = .0005$. There were no differences between the dissociative and

² We have previously reported (Wolf, Lunney, et al., 2012) that the dissociative class did not differ from the high-PTSD group on demographic or trauma-related variables with the exception of race, such that the moderate- and high-PTSD groups included a greater percentage of women who self-reported as White (relative to identifying as a racial minority). Given this, we conducted secondary analyses in which we included race (minority or nonminority) as a covariate of group membership and reran the latent growth model. Results with respect to the effect of the DS on the intercept and growth factors were unchanged from the primary results reported in the text (details available from first author).

high-PTSD classes, Wald's chi-square ($1, n = 235$) = 0.02, $p = .89$. This pattern held at both 3 and 6 months. There were no differences in the number of completed treatment sessions as a function of group membership (moderate-PTSD group: $M = 7.91$, $SD = 3.53$; high-PTSD group: $M = 8.50$, $SD = 2.94$; dissociative group: $M = 8.04$, $SD = 3.33$), $F(2, 281) = 0.78$, $p = .46$, and there was no difference in the number of completed sessions as a function of group membership by treatment type; $F(2, 278) = 0.04$, $p = .96$. There was no difference in the proportion of participants who received any treatment (i.e., one or more sessions) as a function of class membership (moderate group: 93.4%; high group: 94.9%; dissociative group: 94.1%), $\chi^2(2, 284) = 0.19$, $p = .91$, nor was there a Treatment Type \times Class assignment term effect on receiving any treatment, $\chi^2(2, 284) = 0.95$, $p = .62$. There were no group differences in the proportion of participants who completed treatment (moderate group: 70.3%; high group: 74.4%; dissociative group: 68.2%), $\chi^2(2, 284) = 0.77$, $p = .68$, and no interaction between group membership and treatment type on therapy completion, $\chi^2(2, 284) = 0.03$, $p = .99$.

We next tested whether the severity of derealization and depersonalization changed in response to PTSD treatment. Descriptive statistics for each class at each time point as a function of treatment type are listed in Table 3. As with the model for the CAPS, the initial model with the Treatment Type \times Class assignment term yielded good fit, $\chi^2(13, N = 284) = 16.00$, $p = .25$ (RMSEA = .03, 90% CI for RMSEA [$<.001$, .07], CFI = 1.0, TLI = .99, SRMR = .02), but the interaction did not predict the slope or intercept factors (smallest $p = .10$). Given this, we again collapsed across treatment type and reran the latent growth curve model. This model fit the data well, $\chi^2(7, N = 284) = 12.40$, $p = .09$ (RMSEA = .05, 90% CI for RMSEA [$<.001$, .10], CFI = .99, TLI = .99, SRMR = .03), and revealed significant effects of the moderate- and high-PTSD classes (vs. the dissociative class) on the intercept factor (unst γ s = -1.66 and -1.45 , respectively, both p s $< .001$, 95% CIs [-1.77 , -1.55] and [-1.57 , -1.32], respectively; see Figure 2). As expected, the former two groups had lower baseline mean scores on dissociation relative to the dissociative class. In addition, both group variables predicted the




Figure 1. The figure shows the estimated mean CAPS severity scores at pretreatment and follow-up assessments as a function of class assignment. PTSD = posttraumatic stress disorder; CAPS = Clinician Administered PTSD Scale.

Table 2

Percentage of Individuals Who No Longer Met Criteria for PTSD at Follow-Up as a Function of Class Assignment

Follow-up point	% No longer meeting criteria for PTSD			Omnibus comparison χ^2 ($df = 2$)
	1. Moderate PTSD ($n = 98$)	2. High-PTSD ($n = 64$)	3. High-PTSD & dissociation ($n = 73$)	
Month 0	43.88 _a	18.75 _b	17.81 _b	18.19*
Month 3	45.92 _a	24.19 _b	20.83 _b	14.50*
Month 6	50.53 _a	26.15 _b	26.09 _b	14.30*

Note. Sample size listed in the second row reflects the baseline sample size for each group. Sample size at the immediate follow-up assessment (Month 0) was $n = 98$, 64, and 73 for Groups 1, 2, and 3, respectively. Sample size at Month 3 was $n = 98$, 62, and 72 for Groups 1, 2, and 3, respectively. Sample size at Month 6 was $n = 95$, 65, and 69 for Groups 1, 2, and 3, respectively. For each follow-up, percentages sharing the same subscript are not significantly different from each other ($p > .05$). PTSD = posttraumatic stress disorder; df = degrees of freedom. * $p < .001$.

slope factor such that the moderate- and high-PTSD groups had more positive slopes (unst γ s = 0.67 and 0.71, respectively, both p s $< .001$, 95% CIs [0.46, 0.89] and [0.48, 0.94], respectively) relative to the dissociative class, indicating less dissociative symptom decline in these two groups.³ The baseline conditional intercept estimate for the dissociative group was 1.95, 95% CI [1.86, 2.03], $p < .001$, and the average rate of symptom decline for this group was estimated at -0.52 , 95% CI [-0.69 , -0.36], $p < .001$, across the first to last assessment points. The slope and intercept factors were not related to each other, $r = -.04$, $p = .69$. The residual variance for the slope factor was significant ($p = .005$); the residual variance for the intercept factor was not ($p = .213$).

Discussion

To our knowledge, this is the largest study to date to evaluate the effect of the DS on PTSD treatment response and the first to use latent class assignment to the dissociative group as the moderator of treatment outcome. This was also the first time the DS has been shown to moderate overall PTSD treatment response; prior work has supported subtle effects of dissociation on cognitive and narrative therapy responses (Cloitre et al., 2012; Resick et al., 2012) but no overall Dissociation \times Time effects. In our sample of female active duty service members and veterans undergoing a trial of PE versus PCT, the high-PTSD group demonstrated, on average, a 9.75 greater decrease in CAPS scores over the course of the study compared with the dissociative group. The high-PTSD group also had higher baseline PTSD symptoms so that, by 6 months, the high-PTSD group was estimated to have CAPS scores that were just 4.7 points below that of the dissociative class. Thus, although dissociation was associated with lesser treatment response relative to the high-PTSD group, the magnitude of this effect was small and the clinical significance modest. The rate of

³ We also reexamined our latent growth curve model predicting change in dissociative symptoms with race as a covariate of group membership (as Footnote 2). We again found that results with respect to the slope and intercept effects were unchanged from that reported in the text without this covariate (details available from first author).

Table 3

Observed Mean Dissociation Severity Scores (and SDs) by Group and Treatment Type at Each Assessment

Class	Baseline		Month 0		Month 3		Month 6	
	PCT	PE	PCT	PE	PCT	PE	PCT	PE
Moderate PTSD	0.30 (0.33)	0.27 (0.36)	0.47 (0.49)	0.30 (0.43)	0.48 (0.61)	0.38 (0.48)	0.55 (0.71)	0.31 (0.56)
High-PTSD	0.49 (0.36)	0.50 (0.39)	0.71 (0.76)	0.78 (0.78)	0.64 (0.62)	0.74 (0.81)	0.41 (0.55)	0.75 (0.71)
High-PTSD & dissociation	1.92 (0.54)	1.97 (0.48)	1.53 (0.80)	1.47 (0.88)	1.54 (0.88)	1.47 (0.81)	1.42 (0.90)	1.22 (0.97)

Note. PCT = present-centered therapy; PE = prolonged exposure; PTSD = posttraumatic stress disorder. All values are *M* (*SD*).

change did not differ between the dissociative and moderate-PTSD classes. Results also suggested that PTSD treatment yielded marked reductions in dissociative symptoms, consistent with prior work (Cloitre et al., 2012; Hagenaars et al., 2010; Resick et al., 2012). This is notable given that the treatments did not include dissociation-related content. Analyses also suggested that participants with the DS were no more likely to drop out from the trial or to otherwise receive less treatment than those without the DS, regardless of treatment type. This suggests the intervention was tolerable to those with the DS. In contrast to the prevailing clinical perception (Becker et al., 2004), these findings suggest that the DS is not a contraindication for PE, as individuals with the DS can benefit from exposure therapy.

Our findings extend the literature concerning the importance of the DS in the *DSM-5* (American Psychiatric Association, 2013) diagnostic criteria. The DS may prove useful in treatment planning, case conceptualization, and tracking an individual's response to treatment. It would be reasonable to provide psychoeducation regarding dissociation to those with the DS, to ensure that symptoms of dissociation do not endanger an individual's safety, and to develop safety plans as needed. Results also suggest the importance of assessing the DS in future PTSD treatment trials. If the DS is not quantified and its variance separated from that of PTSD, then it is likely that unmeasured variability in the diagnosis will obscure the search for effective treatments.

Results should be considered in light of study limitations. First, this study included only female veterans and active duty service

members undergoing PE and PCT. It is unknown whether results generalize to men, nonveterans, or those engaged in other forms of PTSD treatment. Second, it is possible that excluding individuals with active self-injury and suicidal ideation from the trial may have limited the severity of dissociative symptoms in the sample. Third, analyses focused on DS by Treatment Type interactions on the rate of change are two-way interactions and were likely underpowered. Fourth, results are based on the *DSM-IV* (American Psychiatric Association, 1994) definition of PTSD and it is unclear whether they generalize to the *DSM-5* (American Psychiatric Association, 2013) definition. Finally, given inconsistent findings in the literature and the fact that our significant differences in slope from baseline to 6-month follow-up were somewhat obscured by differences in baseline symptoms between the high-PTSD and dissociative groups, it is important for future work to attempt to replicate these results. Despite these limitations, results provide empirical support for the use of exposure treatment for individuals with the DS of PTSD and highlight the value of the inclusion of the DS in *DSM-5*.

References

- American Psychiatric Association. (1994). *Diagnostic and statistical manual of mental disorders* (4th ed.). Washington, DC: Author.
- American Psychiatric Association. (2013). *Diagnostic and statistical manual of mental disorders* (5th ed.). Arlington, VA: Author.
- Armour, C., Elklit, A., Lauterbach, D., & Elhai, J. D. (2014). The *DSM-5* dissociative-PTSD subtype: Can levels of depression, anxiety, hostility, and sleeping difficulties differentiate between dissociative-PTSD and PTSD in rape and sexual assault victims? *Journal of Anxiety Disorders*, 28, 418–426. <http://dx.doi.org/10.1016/j.janxdis.2013.12.008>
- Becker, C. B., Zayfert, C., & Anderson, E. (2004). A survey of psychologists' attitudes towards and utilization of exposure therapy for PTSD. *Behaviour Research and Therapy*, 42, 277–292. [http://dx.doi.org/10.1016/S0005-7967\(03\)00138-4](http://dx.doi.org/10.1016/S0005-7967(03)00138-4)
- Blake, D. D., Weathers, F. W., Nagy, L. M., Kaloupek, D. G., Gusman, F. D., Charney, D. S., & Keane, T. M. (1995). The development of a clinician-administered PTSD scale. *Journal of Traumatic Stress*, 8, 75–90. <http://dx.doi.org/10.1002/jts.2490080106>
- Briere, J. (1995). *Trauma Symptom Inventory professional manual*. Odessa, FL: Psychological Assessment Resources.
- Cloitre, M., Petkova, E., Wang, J., & Lu Lassell, F. (2012). An examination of the influence of a sequential treatment on the course and impact of dissociation among women with PTSD related to childhood abuse. *Depression and Anxiety*, 29, 709–717. <http://dx.doi.org/10.1002/da.21920>
- Hagenaars, M. A., van Minnen, A., & Hoogduin, K. A. (2010). The impact of dissociation and depression on the efficacy of prolonged exposure treatment for PTSD. *Behaviour Research and Therapy*, 48, 19–27. <http://dx.doi.org/10.1016/j.brat.2009.09.001>




Figure 2. The figure shows the estimated mean dissociation item severity scores at pretreatment and follow-up assessments as a function of class assignment. PTSD = posttraumatic stress disorder; CAPS = Clinician Administered PTSD Scale.

- Halvorsen, J. Ø., Stenmark, H., Neuner, F., & Nordahl, H. M. (2014). Does dissociation moderate treatment outcomes of narrative exposure therapy for PTSD? A secondary analysis from a randomized controlled clinical trial. *Behaviour Research and Therapy*, 57, 21–28. <http://dx.doi.org/10.1016/j.brat.2014.03.010>
- Lanius, R. A., Vermetten, E., Loewenstein, R. J., Brand, B., Schmahl, C., Bremner, J. D., & Spiegel, D. (2010). Emotion modulation in PTSD: Clinical and neurobiological evidence for a dissociative subtype. *The American Journal of Psychiatry*, 167, 640–647. <http://dx.doi.org/10.1176/appi.ajp.2009.09081168>
- Muthén, L. K., & Muthén, B. O. (1998–2012). *Mplus user's guide* (7th ed.). Los Angeles, CA: Author.
- Price, M., Kearns, M., Houry, D., & Rothbaum, B. O. (2014). Emergency department predictors of posttraumatic stress reduction for trauma-exposed individuals with and without an early intervention. *Journal of Consulting and Clinical Psychology*, 82, 336–341. <http://dx.doi.org/10.1037/a0035537>
- Resick, P. A., Suvak, M. K., Johnides, B. D., Mitchell, K. S., & Iverson, K. M. (2012). The impact of dissociation on PTSD treatment with cognitive processing therapy. *Depression and Anxiety*, 29, 718–730. <http://dx.doi.org/10.1002/da.21938>
- Schnurr, P. P., Friedman, M. J., Engel, C. C., Foa, E. B., Shea, M. T., Chow, B. K., . . . Bernardy, N. (2007). Cognitive behavioral therapy for posttraumatic stress disorder in women: A randomized controlled trial. *JAMA: Journal of the American Medical Association*, 297, 820–830. <http://dx.doi.org/10.1001/jama.297.8.820>
- Stein, D. J., Koenen, K. C., Friedman, M. J., Hill, E., McLaughlin, K. A., Petukhova, M., . . . Kessler, R. C. (2013). Dissociation in posttraumatic stress disorder: Evidence from the world mental health surveys. *Biological Psychiatry*, 73, 302–312. <http://dx.doi.org/10.1016/j.biopsych.2012.08.022>
- Steuwe, C., Lanius, R. A., & Frewen, P. A. (2012). Evidence for a dissociative subtype of PTSD by latent profile and confirmatory factor analyses in a civilian sample. *Depression and Anxiety*, 29, 689–700. <http://dx.doi.org/10.1002/da.21944>
- van Minnen, A., Harned, M. S., Zoellner, L., & Mills, K. (2012). Examining potential contraindications for prolonged exposure therapy for PTSD. *European Journal of Psychotraumatology*, 3, 18805.
- Wolf, E. J., Lunney, C. A., Miller, M. W., Resick, P. A., Friedman, M. J., & Schnurr, P. P. (2012). The dissociative subtype of PTSD: A replication and extension. *Depression and Anxiety*, 29, 679–688. <http://dx.doi.org/10.1002/da.21946>
- Wolf, E. J., Miller, M. W., Reardon, A. F., Ryabchenko, K. A., Castillo, D., & Freund, R. (2012). A latent class analysis of dissociation and PTSD: Evidence for a dissociative subtype. *Archives of General Psychiatry*, 69, 698–705. <http://dx.doi.org/10.1001/archgenpsychiatry.2011.1574>

Received December 8, 2014

Revision received April 8, 2015

Accepted May 22, 2015 ■